

BRAIN HEALTH & ALASKA SEAFOOD



Written by Kari Natwick, RDN, LD, IFNCP January 2019



TABLE OF CONTENTS

1. BRAIN HEALTH AND ALASKA SEAFOOD

2. ALZHEIMER'S DISEASE AND DEMENTIA

3. DEPRESSION

4. PERINATAL DEPRESSION


5. VITAMIN D

6. SEAFOOD PROMOTES BRAIN HEALTH

7. SOURCES

BRAIN HEALTH AND ALASKA SEAFOOD

When our brain suffers, so does our quality of life. For many people, brain health is a growing concern as factors in our lives such as stress, toxins, nutrient deficiencies and aging all contribute to changes in our cognition and mood. The impact of lifestyle and diet has been extensively studied on brain health, and we now know that there are multiple interventions that can improve our cognition and reduce risk of dementia, Alzheimer's disease, and mental health disorders. Some of the positive lifestyle changes include increasing physical activity, having healthy social relationships, engaging in mentally stimulating activities, avoiding toxins in our environment, and eating a nutrient-dense whole food diet.



ONE OF THE PRIMARY REASONS SEAFOOD, ESPECIALLY FATTY FISH, CONTRIBUTES TO BRAIN HEALTH

is the fact that it is rich in vital long-chain omega-3 polyunsaturated fatty acids (omega-3's), particularly eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA).

There are many components of a diet to support brain health, and the inclusion of seafood is proving to be essential. DHA is the dominant omega-3 in the brain and is a critical structural component of every cell. DHA has the ability to turn on the growth of new brain cells as well as offer protection for existing brain cells. [9] It also increases neuroplasticity, which is the ability of the brain to connect one brain cell to the next. [9] This allows us to learn, to have new ideas and to heal after an injury to the brain. Omega-3 fatty acids, DHA in particular, also help to decrease inflammation in the brain [9], which can occur in people who have had traumatic brain injuries, spinal cord injuries, Alzheimer's disease, Parkinson's Disease or Multiple Sclerosis.

EPA and DHA are synthesized by the body in small amounts, but not in quantities considered to be enough to support all the vital functions they perform in the body. Couple this with the fact that the majority of Americans do not consume 8-ounces of seafood per week as recommended by the Dietary Guidelines for Americans 2015-2020, and it is easy to see why most people are not getting adequate amounts of these essential fatty acids. [30] Because of this, it is important to focus on the consumption of food sources of omega-3 fatty acids, especially those highest in EPA and DHA.



The best-known sources of EPA and DHA on the planet are high-quality seafood, such as the wild seafood originating from Alaska waters.



ALZHEIMER'S DISEASE AND DEMENTIA

Alzheimer's disease is an irreversible and progressive condition that affects 5.4 million Americans and is the 6th leading cause of death among adults in the U.S. Alzheimer's disease is not fully understood, but it is thought to be the result of a combination of genetic, environmental and lifestyle factors. Alzheimer's disease begins an average of 30 years before the first symptoms are seen, [10] with the risk of developing Alzheimer's disease and other late-life dementia doubling every five years after the age of 65. [1, 31] Due to the constant aging of the population worldwide, the incidence of Alzheimer's disease has increased exponentially. [2] Alzheimer's disease can cost a family both financially and in emotional burden with the chronic stress of having to caretake an adult.

Cognitive decline, which is a response to aging neurons and the decreased speed at which the brain functions, is inevitably linked with the aging process and manifests itself as worsening memory, processing speed and attention.



Consuming as little as one seafood meal per week has been associated with decreasing one's risk for both AD and dementia.

Nutrition interventions, such as increased consumption of fatty fish, and the subsequent increases in EPA and DHA levels in the blood, have proven to decrease the risk of developing dementia and Alzheimer's disease. [5,7, 8, 31] In fact, consuming as little as one seafood meal per week has been associated with decreasing one's risk for both AD and dementia. [4] The reduction in both cognitive decline and Alzheimer's disease which is associated with increased omega-3 intake are attributable to the neurogenerative (production of new brain cells) and anti-inflammatory effects of omega-3 fatty acids. Omega-3 fatty acids also enhance the immune system's ability to clear the brain of beta amyloid plaques, which are one of the hallmarks of Alzheimer's disease. [33, 34] Increases in gray matter, brain volume and cognitive improvements are also evident when individuals have increased levels of EPA and DHA levels in their blood. [3, 6] Low omega-3 fatty acid intake is also a risk factor for the development of cardiovascular disease, diabetes, and hypertension, all of which also increase an individual's risk for the development of cognitive decline and Alzheimer's Disease. [31]

DESPITE THE CLEAR EVIDENCE THAT EATING FISH IS GOOD FOR BRAIN HEALTH, IT'S POSSIBLE THAT OMEGA-3'S INTAKE MAY HAVE AN EVEN GREATER IMPACT ON CERTAIN INDIVIDUALS WHO ARE AT HIGH RISK FOR DEVELOPING ALZHEIMER'S DISEASE AND COGNITIVE DECLINE DUE TO GENETICS.

These are individuals who are carriers of the APOE4 gene. In these individuals, the presence of the APOE4 gene is associated with earlier onset of the symptoms of Alzheimer's Disease, and the protective benefits of DHA from fish appears to be particularly potent, especially when taken during the pre-dementia state of Alzheimer's disease. [31, 32] In one study, consuming one seafood meal per week protected against decline in multiple cognitive domains, especially if an individual was a carrier of the APOE4 gene. [29]

Wild Alaska Seafood

such as salmon, sablefish, halibut, oysters, herring, and many others are some of the greatest sources of brain protective omega-3 fatty acids on the planet.

The evidence is clear that consuming seafood rich in omega-3 fatty acids can protect the brain from cognitive decline and Alzheimer's disease. In order to promote brain health, consuming Alaska seafood twice per week is a proven strategy to decrease the risk of cognitive decline and Alzheimer's disease.

DEPRESSION

Depression is a serious public health problem and is the leading cause of disability worldwide. [15] It affects about 16 million American adults each year, and 1 in 6 adults will have depression in their lifetime. Antidepressants are one of the most commonly used therapeutic drug classes in the United States, and use has increased almost 65% in a 15-year period. 12.7% of people over age 12 are estimated to have taken anti-depressant medication within the past month with approximately 25% of those taking antidepressant medication had done so for 10 years or more. [11]

When it comes to the prevention of depression and reducing symptoms of depression, diet is a well-known factor. [17, 20] Adherence to a nutrient-rich diet that is high in fruits, vegetables, whole grains, olive oil, antioxidants and fatty fish, such as the Mediterranean diet, reduces symptoms of depression. [21, 42] In countries where seafood consumption is higher, such as Japan and Korea, there appears to be lower rates of depression. [16, 18]



Consumption of seafood has been linked to reductions in the rate of post-partum depression [14] as well as depression in general in adults. [13]

The mechanisms by which increased seafood intake reduces rates of depression is still under investigation, but a key link is inflammation. Inflammation that increases levels of proinflammatory cytokines such as interleukins IL-6 and IL-1B and TNF-alpha are known to result in depression and inhibit memory. [39] EPA and DHA, which are found in higher quantities in seafood, and are known for their anti-inflammatory [19], antioxidative, neuroprotective and neurogenesis [22] effects.



EPA AND DHA FROM SEAFOOD

help to protect, restore and rebuild the brain.

These are important factors when protecting against depression as well as other diseases. In addition to diet, other studies have shown that supplementation with EPA (>60% of total EPA and DHA) has shown significant benefits for reducing depression [23], although, again, the mechanism is not clearly defined. EPA is neuroprotective and has anti-inflammatory effects, which are suggestive to drive this effect. [12]

Another key component when it comes to mood and cognition (memory) is a brain chemical called serotonin. Serotonin plays a critical role in brain function by helping information to transmit across the nervous system (neurotransmission) and as a hormone. The majority of serotonin is located in the gut, but it is also located in regions of the brain that have been dubbed "the social brain" as these areas regulate social cognition and decision-making. Serotonin affects a wide-range of cognitive functions, and low levels have been linked to memory problems, low mood (depression), aggression, impulsivity, anxiety, psychopathology and personality disorder. [39]



Wild Alaska salmon contains unique nutritional properties that help to support serotonin production.

Consuming wild Alaska seafood can help boost serotonin in the brain in a variety of ways. First, it is proposed that omega-3 fatty acids support serotonin release into the brain by reducing inflammation in the brain. [39] One of the mechanisms for this is that EPA increases serotonin release by reducing inflammatory properties (E2 series prostaglandins) that inhibit serotonin release. DHA also influences serotonin uptake by making receptors more sensitive, which makes the uptake more effective. This means that when levels of omega-3 fatty acids are low in the blood; the body may not utilize serotonin efficiently. [39]

Next, the amino acid tryptophan is necessary for the production of serotonin. Tryptophan is the sole precursor of serotonin produced in the body, [43] and is therefore necessary for the production of serotonin. However, because the body cannot manufacture tryptophan, it must be acquired through dietary sources. Tryptophan is available in a variety of protein-related food sources, with wild Alaska salmon and halibut being good sources of this amino acid.

It must be noted that despite the importance of serotonin in the brain, it is not an isolating factor in neuropsychiatric disorders as these conditions are multifactorial and influenced by several complex interactions including genetics, nutrition and environment.

When it comes to implementing dietary strategies to reduce the risk of depression and symptoms of depression, including 4-ounces of seafood twice a week as a part of your diet can be beneficial.



Wild Alaska seafood high in EPA and DHA is an important part of many of the proven dietary patterns than enhance mood and reduce depression.



Wild Alaska seafood is a high quality, nutrient dense food that includes the brain supporting nutritional components of EPA, DHA, tryptophan and Vitamin D. Because of this, Alaska seafood is known for its anti-inflammatory and neuroprotective effects and may increase serotonin levels in the brain, all of which are critical elements for brain health.

PERINATAL DEPRESSION

According to the CDC, 1 in 9 women, or 10-20%, experience symptoms of postpartum depression. [35] Symptoms of postpartum depression may include crying more often than usual, feelings of anger, withdrawing from loved ones, feeling numb or disconnected from your baby, and feeling guilty about not being a good mom or doubting your ability to care for your baby. [35]

The amount of omega-3 fatty acids in the blood decline during pregnancy and lactation because the fetus uses omega-3s for the development of its nervous system, and the mother's body uses omega 3's after birth to make breastmilk. [14] Unfortunately, **most pregnant women's diets are inadequate in omega 3's due to low consumption of dietary sources high in omega 3's, such as seafood.** Because of this, blood levels of omega-3 fatty acids in pregnant and breastfeeding woman may not be optimal and can impact not only child development, but may also put a pregnant mother at risk for the development of post-partum depression.

**MOST PREGNANT
WOMEN'S DIETS ARE**



**INADEQUATE IN
OMEGA 3'S**

**DUE TO LOW CONSUMPTION OF
DIETARY SOURCES HIGH IN
OMEGA 3'S, SUCH AS SEAFOOD.**

Research that measures blood levels of omega-3's during pregnancy shows a clear link between low blood levels of omega-3's and increased rates of post-partum depression. [36] Other studies have demonstrated that DHA concentrations in the blood are often significantly lower in postpartum women experiencing depressive symptoms than those who are not. [37,38] In one study, every 1% increase in DHA in the blood was associated with a 59% reduction in the reporting of depressive symptoms in pregnant mothers. [40] It is thought that the benefits of omega-3 fatty acids are related to the reduction in pro-inflammatory cytokines that promote inflammation are elevated during depression.

One of the best food sources of omega-3 fatty acids is seafood; however, pregnant women tend to consume small quantities of fish. One of the reasons women may omit fish in their diet is due to concerns over the mercury content. Methylmercury is a known neurotoxin that accumulates in aquatic food chains with levels that vary in species depending on their size and diet. Mercury can cross the placenta placing a fetus at risk for exposure. Longer living predatory fish such as shark, tilefish, mackerel and swordfish should be avoided due to their high levels of mercury.

Additionally, these foods are excellent sources of selenium, which protects against mercury toxicity.



Table 1
Mercury and Fish Consumption During Pregnancy

Fish to Consume*	Fish to Avoid
Shrimp	Shark
Salmon	Swordfish
Pollock	Mackerel
Scallops	Tilefish

*Have less than 0.05 ppb of mercury per 6-oz serving, except light tuna, which has 1.2 ppb. 1 µg/kg = 1 ppb

Unfortunately, when pregnant women consume small quantities of seafood, their diets are insufficient in omega-3 fatty acids. This dietary omission can place women at risk for developing perinatal depression. Omega-3 fatty acids are well known for their ability to reduce neuroinflammation (inflammation in the brain), which is a key association of perinatal depression. Pregnant women should consume 4-ounces of fatty fish per week twice a week, as is recommended by the 2015 Dietary Guidelines for Americans. Especially important is to include seafood that is high in omega-3 fatty acids and low in methylmercury.



Wild Alaska seafood is an excellent choice since salmon, herring and sablefish are some of the greatest sources of omega 3's on the planet. Because they are wild-caught in the pristine waters of Alaska, they are low in levels of methylmercury and are safe to consume during pregnancy.



[Click here](https://www.fda.gov/food/consumers/eating-fish-what-pregnant-women-and-parents-should-know)¹ for more information on recommendations for pregnant women and children regarding mercury and Alaska seafood.

1) <https://www.fda.gov/food/consumers/eating-fish-what-pregnant-women-and-parents-should-know>

VITAMIN D

Vitamin D3, or cholecalciferol, is a fat-soluble vitamin that functions as a hormone precursor. Vitamin D3 occurs naturally in foods such as fatty fish, as well as in fortified food products such as milk or orange juice. It is also found in small amounts in other foods such as mushrooms exposed to UV light, cheese, egg yolk and beef liver. Vitamin D is also synthesized in the skin after exposure to sunlight or other sources of ultraviolet light. Vitamin D is functional in the body as it enhances the absorption of calcium and phosphorus, which promote bone mineralization and remodeling. Vitamin D also plays a role in neuromuscular function and influences cellular growth and differentiation. Additionally, Vitamin D appears to enhance the secretion and action of insulin.

Throughout most of human history, Vitamin D has been obtained almost exclusively through the skin.

However, due to modern lifestyles, individuals spend less time outdoors and most Western populations are deficient in this key nutrient. [26]

Individuals who are especially at risk for developing Vitamin D deficiency are those who live above 35 degrees in latitude; being elderly or obese; having dark skin; avoiding sunlight exposure through staying indoors or excessive use of sunscreen; having low dietary Vitamin D intake; having a condition that causes malabsorption; breastfed infants; or taking Vitamin D-depleting drugs.[24] Interesting to note is that, with the exception of infants, these are the same populations that are at risk for depression. [28]

Low levels of Vitamin D are often found in individuals who suffer from depression, anxiety and other mental health disorders. [28] The link between Vitamin D and cognitive function is that Vitamin D plays a neuroprotective role. Low levels have also been associated with an increased risk of substantial cognitive decline in several population-based studies. Vitamin D deficiency has also been linked to an array of diseases such as rickets, cardiovascular disease, osteoporosis and cancer. [25]

In order to obtain adequate amounts of Vitamin D, the best ways are adequate exposure to sunshine [27], supplementation or consumption of foods high in Vitamin D. Very few foods naturally contain Vitamin D, so in order to ensure adequate consumption, it is important to include the best sources in your diet. Wild Alaska seafood is an excellent source of Vitamin D. For example, canned **Alaska sockeye salmon provides 720 IU's per 3-ounce portion, meeting 120% of the RDA. Another example is 3-ounces of Alaska sockeye contains 570 IU's of Vitamin D, almost 100% the RDA for an adult.** Alaska seafood is an excellent source of vitamin D.

[Click here](#)¹ to learn more about the nutritional value, including Vitamin D levels, of Alaska seafood.

1) https://uploads.alaskaseafood.org/2018/12/NutritionalValues_Final.pdf

Table 2: Recommended Dietary Allowances (RDAs) for Vitamin D [1]

Age	Male	Female	Pregnancy	Lactation
0–12 months*	400 IU (10 mcg)	400 IU (10 mcg)		
1–13 years	600 IU (15 mcg)	600 IU (15 mcg)		
14–18 years	600 IU (15 mcg)			
19–50 years	600 IU (15 mcg)			
51–70 years	600 IU (15 mcg)	600 IU (15 mcg)		
>70 years	800 IU (20 mcg)	800 IU (20 mcg)		

* Adequate Intake (AI)

SEAFOOD PROMOTES BRAIN HEALTH

The connection between diet and brain health is very clear. Consuming a nutrient-dense diet that contains key nutrients known to reduce one's risk of developing Alzheimer's disease, cognitive decline and depression is essential. Omega-3 fatty acids are one of the key nutrients most people do not consume enough of due to a general decline in the consumption of seafood in Western countries over the past century. Given that the brain is largely made up of omega-3 fatty acids [41], the importance of increasing these fats in our diet deserves greater attention to promote brain. Additionally, blood levels of Vitamin D are often inadequate, which also places people at risk for depression, anxiety and cognitive decline.



Wild Alaska salmon, halibut, and sablefish contain Vitamin D, amino acids such as tryptophan, and the omega-3 fatty acids EPA and DHA, all of which are necessary to supporting mood and cognition. Additionally, wild Alaska seafood is an excellent source of protein, selenium, and B vitamins that are essential to health and well-being.

In order to ensure adequate consumption, consuming 4 ounces of wild Alaska seafood, twice a week is recommended.

Also, aim to consume a daily average of 250 mg of EPA/DHA throughout the week. Last, a whole food nutrient-dense diet based upon the Mediterranean diet, can support the brain and reduce a person's risk of chronic disease. Focusing on these nutrition interventions, with the inclusion of wild Alaska seafood, are important steps toward optimizing brain health and reducing risk of Alzheimer's disease, cognitive decline, and depression.

KEY RECOMMENDATIONS

1. Consume 8 ounces of seafood per week (4 ounces of seafood, twice a week).
2. Focus on consuming fish high in EPA and DHA such as Alaska salmon, sablefish, oysters, halibut, and sardines. Average daily consumption should consist of 250 mg of EPA and DHA per day.
 - 3-ounces of Alaska king salmon contains 1476 mg EPA/DHA
 - 3-ounces Alaska sockeye salmon contains 730 mg EPA/DHA
 - 3-ounces Alaska black cod contains 1543 mg EPA/DHA
 - 3-ounces Alaska halibut contain 201 mg EPA/DHA
 - [Click here](#)¹ for more information on EPA/DHA in Alaska Seafood.
3. Alaska seafood high in EPA and DHA is anti-inflammatory, protects the brain, and boosts serotonin levels.
4. Consume a Mediterranean style diet consisting of nutrient dense whole foods such whole grains, fruits and vegetables, nuts and fish. Avoid processed foods. [Click here](#)² to learn more.

1) https://uploads.alaskaseafood.org/2018/12/NutritionalValues_Final.pdf
2) <https://oldwayspt.org/system/files/atoms/files/StepsforMedDiet.pdf>

SOURCES

1. <https://www.cdc.gov/aging/aginginfo/alzheimers.htm>
2. H. Plassmann et al. *Journal of Consumer Psychology* 22 (2012) 18–36
3. Witte et al. *Cerebral Cortex* November 014; 24:3059-3068
4. Zhang Y et al. *Am J Clin Nutrition*. 2016; 103(2): 330-340
5. *J Clin Med Res*. 2017;9(1):1-9
6. Eriksdotter M et al. *J Alzheimers Dis*. 2015; 48(3): 805-812
7. Grimm et al. *Journal of Lipid Research* 2017; 58(11):2083-2101
8. Klimova et al. *Nutrients* 2018;
9. Cutuili *Current Neuropharmacology* 2017; 15(4): 534-542
10. Kohlstadt *Advancing Medicine with Food and Nutrients, Second Edition*
11. <https://www.cdc.gov/nchs/data/databriefs/db303.pdf>
12. Sublette ME et al. *J. Clin Psychiatry*. 2011; 72(12): 1577-1584
13. Li et al. *J. Epidemiol Community Health*. 2016 Mar;70(3):299-304.
14. Markhus et al. *PLoS One*. 2013 Jul 3;8(7) e67617
15. World Health Organization Health Topics/Depression. [(accessed on 25 October 2017)]; Available online: <http://www.emro.who.int/health-topics/depression/index.html>
16. Wu et al. *J Nutr Health Aging* 2016 Apr;20(4):404-7. doi: 10.1007/s12603-015-0590-0.
17. Li Y. et al. Dietary patterns and depression risk: A meta-analysis. *Psychiatry Res*. 2017;253:373–382. doi: 10.1016/j.psychres.2017.04.020
18. Yang Y1, Je Y2. *Eur J Clin Nutr*. 2018 Aug;72(8):1142-1149. doi: 10.1038/s41430-017-0083-9. Epub 2018 Jan 17.
19. Simopoulos AP1. *J Am Coll Nutr*. 2002 Dec;21(6):495-505.
20. Bountziouka V et al. *J Aging Health* 2009 Sep;21(6):864-80. doi: 10.1177/0898264309340693. Epub 2009 Jul 8.
21. Sanchez-Villegas A et al. *Public Health Nutr*. 2006 Dec;9(8A):1104-9.
22. Kang JX, Gleason ED. *CNS Neurol Disord Drug Targets* 2013 Jun;12(4):460-5.
23. Sublette ME et al. *J Clin Psychiatry* 2011 Dec;72(12):1577-84. doi: 10.4088/JCP.10m06634. Epub 2011 Sep 6.
24. Gaby A. (2011) *Nutritional Medicine* Fritz Perlberg Publishing, Concord, NH
25. Wallis DE, Penckofer S, Sizemore GW *Circulation*. 2008 Sep 30;118(14): 1476-85
26. Berk, Sanders, et al *Med Hypotheses* 2007; 69(6): 1316-9. Epub 2007 May 11
27. <https://ods.od.nih.gov/factsheets/VitaminD-HealthProfessional/>
28. Penckofer S. et al. *Issues Ment Health Nurs* 2010 Jun; 31(6): 385-93
29. Rest O. et al. *Neurology* 2016 May 31; 86(22): 2063-2070
30. Jahns L. et al. *Nutrients* 2014 Dec; 6(12): 6060-6075
31. Cole G. et al. *Prostaglandins Leukot Essent Fatty Acids* 2009 Aug-Sep; 81(0): 213-221
32. Patrick RP *FASEB J* 2018 Oct 5: fj201801412R
33. Fiala M. et al. *FASEB* 2015 March: 2681-2689
34. Flala M. et al. *Journal of Alzheimer's Disease* 2015: 48(2): 293-301
35. <https://www.cdc.gov/reproductivehealth/depression/index.htm>
36. Golding J, Steer C, et al. *Epidemiology* 2009; 20: 598-603
37. Sontrop J, Avison, W.R., et al *Paediatric and Perinatal Epidemiology* 2008; 22(4): 389-399
38. Sontrop J, et al, *Epidemiology* 2009; 20(4); 598-603
39. Patrick R., Ames B *FASEB* 2015 (29): 2207-2222
40. Makrides M et al. *Asia Pac J Clin Nutr*. 2003; 12(Suppl):S37
41. Vourre JM. *J Nutr Health Aging* 2004; 8(3): 163-174
42. Jacka F, O'Neil A. et al *BMC Medicine* 2017 15:23
43. Richards DM, Dawes MA, et al *Int J Tryptophan Res* 2009 Mar 23; 2(): 45-60